

There is no first quantization – except in the de Broglie–Bohm interpretation

Hrvoje Nikolić

Theoretical Physics Division, Rudjer Bošković Institute,
P.O.B. 180, HR-10002 Zagreb, Croatia
hrvoje@thphys.irb.hr

February 1, 2008

Abstract

The relativistic effects of the integer-spin quantum field theory imply that the wave functions describing a fixed number of particles do not admit the usual probabilistic interpretation. Among several most popular interpretations of quantum mechanics applied to first quantization, the only interpretation for which this fact does not lead to a serious problem, and therefore the only consistent interpretation of first quantization, is the de Broglie–Bohm interpretation.

PACS: 03.65.Ta; 03.65.Pm; 03.70.+k

Keywords: De Broglie–Bohm interpretation; Relativistic quantum mechanics; Quantum field theory

Recently, Zeh argued that “there is no first quantization” [1], i.e. that true quantum theory is “second” quantization, known also under the name *quantum field theory* (QFT). According to this paradigm, particles appear as small localized objects not because the fundamental objects are quantized pointlike objects called particles, but rather due to decoherence caused by local interactions between fields.

In this Letter we present further arguments against first quantization, by emphasizing that the usual postulates and interpretations of quantum mechanics (QM) applied to first quantization are inconsistent with relativistic QFT. Specifically, we consider the Copenhagen interpretation, the minimal probabilistic interpretation, the consistent-history interpretation, the Everett interpretation and the de Broglie–Bohm interpretation, and emphasize that all these interpretations except the last one require one of the basic postulates of QM: the existence of a quantity that can be naturally interpreted as a probability. On the other hand, from relativistic QFT it follows that this postulate cannot be applied to wave functions that describe a fixed number of particles. Therefore, as far as *only first* quantization is considered, only two points of view seem to be consistent. One consistent point of view is that decoherence is the best we can say about the problem of measurement. (The decoherence paradigm [2, 3, 4, 5] is not an interpretation at all; decoherence is a physical phenomenon for which there are no doubts that it occurs in nature.) The other consistent point of view is the de Broglie–Bohm interpretation [6, 7, 8, 9, 10] formulated in a form in which both the fields and the particle trajectories are physical entities that objectively exist [11, 12].

Before discussing various interpretations of QM, let us first state a few basic phenomenological rules of QM that are experimentally confirmed and thus should be consistent with any consistent interpretation, at least in the cases for which the experimental confirmation exists. Let X denote some set of the degrees of freedom, say positions of n particles. Assume that a wave function $\psi(X, t)$ can be rewritten in terms of some functions $\psi_i(X, t)$ as a linear combination

$$\psi(X, t) = \sum_i c_i(t) \psi_i(X, t), \quad (1)$$

where $c_i(t)$ are complex coefficients. Assume also that a unitary scalar product $(\ , \)$ of wave functions can be defined such that

$$(\psi_i, \psi_j) = \delta_{ij}, \quad (2)$$

independently of time t . Assume further that there exists a hermitian operator $\hat{A}$ acting on the X -space, such that

$$\hat{A}\psi_i(X, t) = a_i(t)\psi_i(X, t), \quad (3)$$

where $a_i(t)$ are real functions. If all these assumptions are fulfilled, then we can apply the following rules:

1. $\hat{A}$ represents an observable and the probability of finding the system at time t to have the value $a_i(t)$ of the observable $\hat{A}$ is equal to $|c_i(t)|^2$.
2. If the system is found to have the value a_i of the observable $\hat{A}$ at time t , then the probabilities at later times should be calculated by replacing ψ with ψ_i .

Note that, by stating the rules above, we do not claim that the mentioned assumptions *are* fulfilled. Instead, we merely state what we may conclude *if* they are fulfilled. These rules say nothing about the case in which the mentioned assumptions are not fulfilled.

Let us now shortly discuss particular interpretations. (A more detailed review of all most popular interpretations of QM can be found in [13].) All these interpretations are attempts to formulate *fundamental principles valid in all physically possible cases* from which the phenomenological rules above can be derived.

We start with the Copenhagen interpretation. The claim that $|c_i(t)|^2$ represent probabilities is one of its fundamental postulates. This postulate implies that a unitary scalar product with the property (2) *must* exist, which is related to the fact that probabilities cannot be negative and that the sum $\sum_i |c_i(t)|^2$ of all probabilities, at each time, must be equal to 1. Another fundamental postulate says that, in the process of measurement, the wave function ψ “collapses” into ψ_i , i.e. after the measurement the wave function becomes *equal* to ψ_i .

The minimal probabilistic interpretation avoids the introduction of a notorious wave-function-collapse postulate [13]. However, similarly to the Copenhagen interpretation, it postulates that $|c_i(t)|^2$ represent probabilities, which implies that a unitary scalar product with the property (2) *must* exist.

The basic objects in the consistent-history interpretation [14, 15, 16] are sequences of projection operators at different times – called histories. The consistent-history interpretation generalizes the usual rules for calculating probabilities in QM by postulating the rules for calculating the probabilities of histories. Therefore, the consistent-history interpretation also relies on the assumption of the consistency of the probabilistic interpretation.

The Everett relative-state interpretation [17, 18] (some variants of which are also known under the name “many-worlds interpretation” [19]) avoids a postulation of an *a priori* probabilistic interpretation. The only fundamental postulate in this interpretation is the usual causal evolution of the wave function. Therefore, this is a completely deterministic theory. However, this is

not a deterministic theory of observables. Instead, this is merely a deterministic theory of wave functions. Therefore, in this interpretation, the explanation of the rule that $|c_i(t)|^2$ represent the probabilities is a very controversial subject [19]. In order to explain it, some researchers use rather indirect and vague arguments [17, 20] or base their arguments on additional assumptions [21, 22, 23], while others conclude (in our opinion, correctly) that the probabilistic rule cannot be explained in the framework of the Everett interpretation [24, 25, 26]. What is even more important for our discussion, all existing arguments for the probabilistic rule are based on the assumptions that we mentioned before expressing this rule. In the framework of the Everett interpretation, it is not clear at all how one might, even in principle, calculate the probabilities when these assumptions are not fulfilled.

The de Broglie–Bohm (dBB) interpretation [6, 7, 8, 9, 10] is a deterministic nonlocal hidden-variable interpretation. In contrast with the Everett interpretation, it is not only a deterministic theory of wave functions, but also a deterministic theory of observables. In the case of nonrelativistic first quantization, its basic postulate is that the particle velocities are given by (we take $\hbar = 1$)

$$V = \frac{1}{m} \partial_X S(X, t), \quad (4)$$

where m is the mass of a particle and S is the phase of the wave function $\psi(X, t) = R(X, t)e^{iS(X, t)}$. No additional assumption is required for the properties of $\psi(X, t)$. Instead, it is rather a *prediction* of this theory that *if* ψ satisfies the usual nonrelativistic Schrödinger equation of the form

$$-\frac{\nabla^2 \psi}{2m} + U(X)\psi = -i\partial_t \psi, \quad (5)$$

which provides that

$$\frac{d}{dt} \int d^{3n}X |\psi(X, t)|^2 = 0, \quad (6)$$

then the statistical distribution of particles ρ is given by

$$\rho(X, t) = |\psi(X, t)|^2 \quad (7)$$

for every t , provided that at an initial time t_0

$$\rho(X, t_0) = |\psi(X, t_0)|^2. \quad (8)$$

Eq. (8) is also often postulated as one of the fundamental postulates of the dBB interpretation, but this postulate is not really necessary. Instead, one can show that if (5) is satisfied and the interactions are sufficiently complex, then the distribution (7) is the most probable [27, 28] (see also [29, 30]).

It is, of course, well known that the probability distribution (7) is a special case of the general quantum mechanical rule for calculating probabilities explained above. This is most easily seen by writing (1) as

$$\psi(X, t) = \int d^{3n}X' \psi(X', t) \delta^{3n}(X' - X). \quad (9)$$

The scalar product is

$$(\psi_i, \psi_j) = \int d^{3n}X \psi_i^*(X, t) \psi_j(X, t), \quad (10)$$

which is time independent, owing to (5). Since the quantity (ψ_i, ψ_i) is positive definite, the functions ψ_i can be chosen such that (2) is satisfied. Therefore, we see that the dBB interpretation *explains*, rather than postulates, the rules for calculating probabilities that particles are at certain positions.

Now consider *relativistic* QM. In a relativistic generalization of the Schrödinger equation, the wave function of a single free particle with an integer spin satisfies the Klein-Gordon equation (we take $\hbar = c = 1$)

$$(\partial_t^2 - \nabla^2 + m^2)\psi(\mathbf{x}, t) = 0. \quad (11)$$

It is well known that, owing to the second time derivative in (11), ψ does not satisfy (6). Therefore, the quantity $|\psi|^2$ cannot be interpreted as a probability density. If, in (6), $|\psi|^2$ is replaced by $j_0 \equiv i(\psi^* \partial_t \psi - \psi \partial_t \psi^*)$, then such a modified Eq. (6) is satisfied. However, the density j_0 is not positive definite, so it cannot be interpreted as a probability density either. The time-independent scalar product is

$$(\psi_i, \psi_j) = i \int d^3x (\psi_i^* \partial_t \psi_j - \psi_j \partial_t \psi_i^*). \quad (12)$$

The quantity (ψ_i, ψ_i) is not positive definite, which is in contradiction with the assumption (2). A consistent interpretation of ψ is given by second quantization, i.e. QFT (see e.g. [31]). In QFT, (11) becomes an equation for the operator $\hat{\psi}$. The quantity ψ is not a wave function but an observable attributed to the operator $\hat{\psi}$. The wave functional $\Psi[\psi(\mathbf{x}), t]$ determines probabilities of these observables according to the usual quantum mechanical rule. An n -particle wave function can be defined as a matrix element of the form $\langle 0 | \hat{\psi}(\mathbf{x}_1, t) \cdots \hat{\psi}(\mathbf{x}_n, t) | \Psi \rangle$. From Eq. (11) applied to $\hat{\psi}$ one can see that wave functions satisfy multiparticle generalizations of (11), which are equations with the second time derivatives.

Now we see that interpretations of first quantization that contain the probabilistic interpretation as one of its fundamental postulates are not consistent. One can propose that such interpretations are valid only in the case when the relativistic effects are negligible, but in that case, such interpretations cannot be viewed as fundamental principles. Instead, they should be consequences of some more fundamental principles. If one postulates that the probabilistic interpretation is fundamental only at the level of wave functionals of QFT, then, even in the nonrelativistic limit, it does *not* follow at all from this interpretation that the probabilistic interpretation can be applied at the level of n -particle wave functions.

There are 3 possible ways to cope with this problem. The first approach is to take the rules of first quantization as unfundamental phenomenological rules valid only under restricted conditions, without wondering about the true reasons why these rules are valid. The second approach, probably the most popular one at the moment, is to view these rules as consequences of the decoherence at the level of true quantum degrees of freedom – fields [1]. This approach is so popular because it explains, rather than postulates, how the quantum interaction with a sufficiently complex environment forces the state of a system to “collapse” to a state that corresponds to a diagonal (i.e. classical) density matrix, using only the properties of solutions of the Schrödinger equation or any of its generalizations. However, as admitted by most experts for decoherence, it does not explain the “collapse” of the wave function. In other words, decoherence explains why we can use classical statistics to statistically describe the outcome of quantum measurements, but it does not explain why and how the measured observables take definite values.

The third possible approach is the dBB interpretation. Actually, it is important to emphasize that the usual form of the dBB interpretation [6, 7, 8, 9, 10] applies the dBB interpretation either to particles or to fields, not to both. In fact, it is often interpreted as a theory in which, at a fundamental level, bosons are fields and fermions are particles. However, in a recent variant of the dBB interpretation both the fields and the particle trajectories simultaneously exist as objective entities [11, 12]. (A similar variant of the dBB interpretation is presented also in [32], but with an important difference that, contrary to the variant in [11, 12], it is not

relativistic and completely deterministic.) Therefore, the dBB interpretation can be formulated such that the usual rules for calculating probabilities in nonrelativistic first quantization can be *derived* from more fundamental principles that respect the relativistic effects. For a known wave function, in general, it is not so easy to calculate the probabilities of particle positions, except in the nonrelativistic limit. However, these probabilities can be calculated in principle from the deterministic theory of particle trajectories determined by a relativistic generalization of (4). In this sense, the relativistic generalization of the dBB interpretation of QM is able to give testable predictions on certain relativistic quantum phenomena on which the conventional theory does not give any prediction at all.

We also want to emphasize that two consistent points of view: the decoherence paradigm and the dBB interpretation – are not two competitive points of view. Instead, the decoherence is an important ingredient of the dBB interpretation, without which the dBB interpretation cannot explain the usual rules for calculating probabilities. To see why, note first that the dBB interpretation predicts the distribution of momenta to be given by

$$\rho(P, t) = \int d^{3n}X |\psi(X, t)|^2 \delta^{3n}(P - \partial_X S(X, t)), \quad (13)$$

which, in general, is *not* equal to the quantum mechanical distribution $|\tilde{\psi}(P, t)|^2$ [where $\tilde{\psi}(P, t)$ is the Fourier transform of $\psi(X, t)$]. However, this problem resolves when the general theory of quantum measurements in the dBB interpretation [6, 7, 10] is taken into account. This theory of measurements has much in common with the theory of quantum measurements based on decoherence, which is perhaps the clearest from the presentation in [10]. In a way, the dBB interpretation can be viewed as a completion of the theory of measurements based on decoherence, because the dBB interpretation uses the basic ideas of decoherence, but at the same time provides an answer to the question that decoherence by itself cannot answer, namely why and how observables take definite values.

By taking into account the relativistic effects in free QFT, in this Letter we have discredited all discussed interpretations of first quantization, except the dBB interpretation. Although it is not the aim of this Letter to discuss in detail the effects of field interactions, we note that the *linearity* of the Schrödinger equation is also one of the fundamental postulates in some interpretations of QM. On the other hand, the wave functions that result from *interacting* QFT do not satisfy linear equations. These nonlinearities do not lead to any fundamental problem in the dBB interpretation of first quantization [11, 12].

To conclude, we repeat that among several most popular interpretations of QM we discussed, only the dBB interpretation can be applied to first quantization, because other interpretations rely on assumptions that are not really satisfied. Therefore, if any of these interpretations of first quantization is consistent, it can only be the dBB interpretation. Of course, it is possible that some other interpretations of first quantization, not considered in this paper, are also consistent. In particular, it is certainly possible that some other deterministic hidden-variable interpretations, more complicated than the dBB interpretation, are consistent. However, it was not possible to discuss all existing interpretations, so the discussion of this Letter is restricted to the most popular ones.

This work was supported by the Ministry of Science and Technology of the Republic of Croatia under Contract No. 0098002.

References

- [1] H. D. Zeh, Phys. Lett. A 309 (2003) 329.

- [2] H. D. Zeh, *Found. Phys.* 1 (1970) 69.
- [3] E. Joos, H. D. Zeh, *Z. Phys. B* 59 (1985) 223.
- [4] W. H. Zurek, *Phys. Today* 44 (1991) 36.
- [5] R. Omnès, *Phys. Rev. A* 56 (1997) 3383.
- [6] D. Bohm, *Phys. Rev.* 85 (1952) 166, 180.
- [7] D. Bohm, B. J. Hiley, P. N. Kaloyerou, *Phys. Rep.* 144 (1987) 323.
- [8] D. Bohm, B. J. Hiley, *Phys. Rep.* 144 (1987) 349.
- [9] P. R. Holland, *Phys. Rep.* 224 (1993) 95.
- [10] P. R. Holland, *The Quantum Theory of Motion*, Cambridge University Press, Cambridge, 1993.
- [11] H. Nikolić, [quant-ph/0208185](#).
- [12] H. Nikolić, [quant-ph/0302152](#).
- [13] F. Laloë, *Am. J. Phys.* 69 (2001) 655.
- [14] R. B. Griffiths, *J. Stat. Phys.* 36 (1984) 219.
- [15] R. Omnès, *J. Stat. Phys.* 53 (1988) 893.
- [16] M. Gell-Mann, J. Hartle, *Phys. Rev. D* 47 (1993) 3345.
- [17] H. Everett, *Rev. Mod. Phys.* 29 (1957) 454.
- [18] J. A. Wheeler, *Rev. Mod. Phys.* 29 (1957) 463.
- [19] Y. Ben-Dov, *Am. J. Phys.* 58 (1990) 829.
- [20] M. Tegmark, *Fortsch. Phys.* 46 (1998) 855.
- [21] D. Deutsch, *Proc. Roy. Soc. Lond. A* 455 (1999) 3129.
- [22] D. Wallace, [quant-ph/0211104](#); [quant-ph/0303050](#).
- [23] S. Saunders, [quant-ph/0211138](#).
- [24] L. E. Ballentine, *Found. Phys.* 3 (1973) 229.
- [25] A. Kent, *Int. J. Mod. Phys. A* 5 (1990) 1745; [gr-qc/9703089](#).
- [26] H. Barnum, C. Caves, J. Finkelstein, C. Fuchs, R. Schack, *Proc. Roy. Soc. Lond. A* 456 (2000) 1175.
- [27] D. Bohm, *Phys. Rev.* 89 (1953) 458.
- [28] A. Valentini, *Phys. Lett. A* 156 (1991) 5.
- [29] D. Dürr, S. Goldstein, N. Zanghí, *J. Stat. Phys.* 67 (1992) 843.
- [30] D. Dürr, S. Goldstein, N. Zanghí, *Phys. Lett. A* 172 (1992) 6.

- [31] J. D. Bjorken, S. D. Drell, Relativistic Quantum Fields, McGraw-Hill, New York, 1965.
- [32] D. Dürr, S. Goldstein, R. Tumulka, N. Zanghi, J. Phys. A 36 (2003) 4143; quant-ph/0303156.